

THEIN!

OFFICE OF NAVAL RESEARCH
Contract #NR 053-683
Technical Report No. 1



Catalytic Oxidation of Phosphines by
Transition Metal-Activated Carbon Dioxide

by

Kenneth M. Nicholas

Prepared for Publication in the

Journal of Organometallic Chemistry



Boston College

Department of Chemistry

Chestnut Hill, MA. 02167

November 19, 1979

Reproduction in whole or in part is permitted for any purpose of the United States Government

*This document has been approved for public release and sale; its distribution is unlimited

DDC FILE COPY

*	
SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)	READ INSTRUCTIONS
REPORT DOCUMENTATION PAGE 1. REPORT NUMBER 12. GOVT ACCESSION NO.	BEFORE COMPLETING FORM 3. RECIPIENT'S CATALOG NUMBER
1 (one)	
A TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED
Catalytic Oxidation of Phosphines by	interim
Transition Metal-Activated Carbon Dioxide	PERFORMING ORG. REPORT NUMBER
7. AUTHOR(*)	B. CONTRACT OR GRANT NUMBER(9)
Kenneth M. Nicholas	NR 053 683
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Department of Chemistry Boston College, Chestnut Hill MA 02167	
11. CONTROLLING OFFICE NAME AND ADDRESS	12_DEPORT DATE
Office of Naval Research (/// 1	November of Pages
Arlington, VA 22217	4
14. MONITORING AGENCY NAME & ADDRESS(IT distrent from Controlling Pilice)	15. SECURITY CLASS. (of this report)
(/2/6/	unclassified
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
This document has been approved for public release and sale; its distribution is unlimited.	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)	
same as above	
submitted for publication in the Journal of Organometallic Chemistry	
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)	
catalytic oxidation, carbon dioxide activation, phosphine	
oxidation oxidation, carbon dioxide activation, phosphine	
20, ABSTRACT (Continue on reverse side if necessary and identify by block number)	
The complexes (PhiP) iRhCl and (cyclooctene) iRhCl is have been found to catalyze the oxidation of phosphines by carbon dioxide in refluxing decalin. The rate of oxidation increases in the order PPhi PBuPhi PEti.	

DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE 5/N 0102-LF-014-6601

الروز مهجها

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

403 740 JM

80 1 11 00

CATALYTIC OXIDATION OF PHOSPHINES BY TRANSITION METAL-ACTIVATED CARBON DIOXIDE

K.M. NICHOLAS

Department of Chemistry

Boston College, Chestnut Hill, MA 02167 U.S.A.

Summary

The complexes (Ph₃P)₃RhCl and [(cyclooctene)₂RhCl]₂ have been found to catalyze the oxidation of phosphines by carbon dioxide in refluxing decalin. The rate of oxidation increases in the order PPh₃ < PBuPh₂ < PEt₃.

Accession For

NTIS GNANI
DDC TAB
Uncommounced
Juntification

Ey______
Pisteibution/
/veilability Sea

Availability Sea

Special

80 1

As the world's petroleum reserves dwindle, there is an intensive search underway for alternative fuels and organic feedstocks. This has stimulated considerable interest in the potential activation of carbon dioxide by means of transition metal complexes¹. Although a few discreet CO₂ complexes have been reported²⁻⁷, in several cases, reactions of carbon dioxide with metal complexes have yielded carbonyl and carbonate or oxide species - products of CO₂ disproportionation⁷⁻¹¹. In a few instances, e.g. in reactions of CO₂ with (C₂H₄)Mo(diphos)₂¹² and (R₃P)₃RhCl¹³, stoichiometric O-atom transfer to phsophine apparently occurs, producing phosphine oxide, either free or coordinated. We now report that under appropriate conditions, phosphines can be oxidized by carbon dioxide <u>catalytically</u> in the presence of [(cyclooctene)₂RhCl]₂ or (Ph₃P)₃RhCl (eq. 1).

$$CO_2 + PR_3 \xrightarrow{L_3Rh(I)X} OPR_3 + CO$$
 (eq. 1)

When a decalin solution of triphenylphosphine containing one mole % of $(Ph_3P)RhCl$ or $[(C_8H_{14})_2RhCl]_2$ is refluxed (ca. $185^{\circ}C)$ under an atmosphere of purified CO_2^{*} , triphenylphosphine oxide is slowly produced (turnover number ca. 20/day) as indicated by the growth of IR bands at 1210, 1115, and $720cm^{-1**}$. After several days, high yields of Ph_3PO (m.p. $148-151^{\circ}$, $1it^{16}$ 150-152) could be obtained upon cooling the mixture. The co-production of CO was confirmed by the gradual formation of a MnO_2/Ag precipitate when


^{*&}quot;Bone dry" grade ${\rm CO_2}$ was passed first through a ${\rm Cr(II)/HClO_4}$ solution, then through a dry ice trap, and finally over 3A molecular sieves.

^{**}No reaction was observed in the absence of $(PPh_3)_3RhCl$ or under a nitrogen atmosphere.

the exit gases were passed through an aqueous $KMnO_4/AgNO_3$ solution 17.

Other phosphines are also oxidized under these conditions and qualitatively, the rate of reaction parallels the phosphine basicity. Thus, while the oxidation of PPh₃ was complete only after 7-8 days, PBuPh₂ was totally converted to OPBuPh₂ after 3 days, and PEt₃ to OPEt₃ in less than 44 hrs.

We do not as yet have any direct mechanistic information on these reactions and so any proposed mechanisms at present are best considered somewhat speculative. However, in view of the reported conversion of $(R_3P)_2Rh(CO_2)Cl$ to $(R_3P)(R_3PO)Rh(CO)Cl^{13}$ and our observation of the lesser catalytic activity of $(PPh_3)_2Rh(CO)Cl$ compared to $(PPh_3)_3RhCl$ (precluding the intermediacy of the former), we suggest the following scheme as a reasonable working hypothesis.



Kinetic studies are in progress as are efforts to establish the scope of transition metal-mediated CO₂ oxidations with other catalysts and substrates.

ACKNOWLEDGEMENT. The author is grateful for support provided by the Office of Naval Research as well as loans of rhodium chloride from Matthey Bishop and Englehardt Industries.

References

- For a recent review see: M.E. Volpin and L.S. Kolominikov, Organometallic Reactions, vol. 5, 313 (1975);
- M. Aresta, C.F. Nobile, V.G. Albano, E. Forni, M. Manassero, Chem. Comm., 636 (1975).
- 3. M. Aresta and C.F. Nobile, J. Chem. Soc. Dalton, 708 (1977).
- 4. I.S. Kolminikov, A.I. Gusev, G.G. Alexandrov, T.S. Lobeeva, Y.T. Struckov, and M.E. Volpin, J. Organometal. Chem., 59, 349 (1973).
- 5. B.R. Flynn and L. Vaska, J. Am. Chem. Soc., 95, 508 (1973).
- 6. G. Fachinetti, C. Floriani, and P. Zanazzi, <u>ibid</u>, <u>100</u>, 7403 (1978).
- 7. T. Herskovitz, J. Am. Chem. Soc., 99, 2391 (1977).
- 8. J. Chatt, M. Kubota, G. Leigh, F.C. March, R. Mason, and D.J. Yarrow, Chem. Comm., 1033 (1974).
- 9. S. Krogsrud, S. Komiya, T. Ito, J.A. Ibers, and A. Yamamoto, Inorg. Chem., 15, 2798 (1976).
- G. Fachinetti, C. Floriani, A. Chiesi-Villa, and C. Guastini,
 J. Am. Chem. Soc., 101, 1767 (1979).
- 11. G.O. Evans, W.F. Walter, D.R. Mills, and C.A. Streit, J. Organometal. Chem., 144, C34 (1978).
- 12. T. Ito and A. Yamamoto, J. Chem. Soc. Dalton, 1398 (1975).
- 13. M. Aresta and L.F. Nobile, Inorg. Chim. Acta., 24, L 49 (1977).
- 14. G.M. Kosolapoff, J. Am. Chem. Soc., 64, 2982 (1942).
- 15. "The Chemists' Companion," A.J. Gordon and R.A. Ford, Wiley and Sons, N.Y., p. 440 (1972).